

## REVIEW

### Efflux-mediated multiresistance in Gram-negative bacteria

K. Poole

Department of Microbiology and Immunology, Queen's University, Kingston, Ontario, Canada

#### ABSTRACT

Multiresistance in Gram-negative pathogens, particularly *Pseudomonas aeruginosa*, *Stenotrophomonas maltophilia*, *Acinetobacter* spp. and the Enterobacteriaceae, is a significant problem in medicine today. While multiple mechanisms often contribute to multiresistance, a broadly distributed family of three-component multidrug efflux systems is an increasingly recognised determinant of both intrinsic and acquired multiresistance in these organisms. Homologues of these efflux systems are also readily identifiable in the genome sequences of a wide range of Gram-negative organisms, pathogens and non-pathogens alike, where they probably promote efflux-mediated resistance to multiple antimicrobials. Significantly, these systems often accommodate biocides, raising the spectre of biocide-mediated selection of multiresistance in Gram-negative pathogens. While there is some debate as to the natural function of these efflux systems, only some of which are inducible by their antimicrobial substrates, their contribution to resistance in a variety of pathogens nonetheless makes them reasonable targets for therapeutic intervention. Indeed, given the incredible chemical diversity of substrates accommodated by these efflux systems, it is likely that many novel or yet to be discovered antimicrobials will themselves be efflux substrates and, as such, efflux inhibitors may become an important component of Gram-negative antimicrobial therapy.

**Keywords** Efflux, gram-negative, multidrug, multiresistance, resistance

**Accepted:** 30 March 2003

*Clin Microbiol Infect* 2004; 10: 12–26

#### INTRODUCTION

While it is clear that bacterial resistance to antimicrobials pre-dates their clinical use, medical and agricultural practices of the past 50 years or more have promoted resistance development and spread in both human and animal pathogens, thereby compromising effective chemotherapy of infectious diseases. While much attention is focused currently on Gram-positive and mycobacterial pathogens, Gram-negative bacteria, particularly those exhibiting a multiresistance phenotype, can still pose a serious infectious disease threat [1,2]. Resistance to multiple antibiotics is increasingly seen in the Enterobacteriaceae, *Pseudomonas aeruginosa*, *Stenotrophomonas maltophilia* and *Acinetobacter* spp. [3]. Significantly,

infections caused by such multiresistant Gram-negative pathogens are associated with excess morbidity and mortality [4–8].

Resistance to antibiotics occurs typically as a result of drug inactivation or modification, target alteration, or reduced accumulation associated with decreased permeability and/or increased efflux [9]. It may be an intrinsic feature of an organism, or may result from mutation or the acquisition of exogenous resistance genes [10,11]. Thus, multiresistance typically results from the accumulation of multiple mutations and/or resistance genes (e.g., on integrons [12]), but specific growth states (e.g., biofilms) and single mutations (e.g., impacting on outer-membrane permeability or the expression of broadly specific multidrug efflux systems) can also promote multiresistance [13]. This review highlights recent advances in our understanding of efflux-mediated multiresistance in Gram-negative bacteria, including updates on the distribution of efflux systems in these organisms, their substrate profiles, and their contribution to antimicrobial resistance in clinical

Corresponding author and reprint requests: K. Poole, Department of Microbiology and Immunology, Queen's University, Kingston, Ontario, Canada, K7L 3N6  
Tel: +1 613 533 6677  
Fax: +1 613 533 6796  
E-mail: [poolek@post.queensu.ca](mailto:poolek@post.queensu.ca)

strains. The ongoing discussions concerning the natural function of these efflux systems and their possible involvement in biocide-mediated selection of antibiotic-resistant strains will also be highlighted. The interested reader is referred to other recent reviews for a more comprehensive treatment of the subject [13,14].

## MULTIDRUG EFFLUX

While efflux as a mechanism of resistance to individual agents has been known for some time, multidrug efflux systems have only recently been identified and appreciated as significant determinants of resistance [14,15]. Chromosomally encoded and broadly distributed in Gram-negative bacteria [16], these systems play an important role in intrinsic and acquired multiresistance [13,14]. Bacterial efflux systems capable of accommodating multiple antimicrobials fall into five classes: the major facilitator superfamily (MFS); the ATP-binding cassette (ABC) family; the resistance-nodulation-division (RND) family; the small multidrug resistance (SMR) family (a member of the much larger drug/metabolite transporter (DMT) superfamily); and the multidrug and toxic compound extrusion (MATE) family [17]. Although not unique to Gram-negative bacteria, RND family transporters are most commonly found in such organisms and typically operate as part of a tripartite system that includes a periplasmic membrane fusion protein (MFP) and an outer-membrane protein (now called outer-membrane factor (OMF)) [18], an organisation also seen on occasion with MFS (e.g., EmrAB-TolC; see below) and ABC (e.g., the macolide-specific MacAB-TolC efflux system [19]) transporters (Fig. 1). Members of all but the ABC family (whose members hydrolyse ATP to drive drug efflux) function as secondary transporters, catalysing drug-ion ( $H^+$  or  $Na^+$ ) antiport (Fig. 1).

## RND FAMILY OF MULTIDRUG EFFLUX SYSTEMS

The most relevant multidrug efflux systems in terms of resistance to clinically important agents (including antibiotics and biocides) are members of the RND family [13] (Table 1), although members of the MFS, MATE and SMR families also show a limited ability to promote resistance to some biocides and antibiotics (Table 2). Originally iden-




Fig. 1. The organisation and operation of antimicrobial efflux pumps of Gram-negative bacteria. OM, outer membrane; PP, periplasmic space; CM, cytoplasmic membrane.

tified as determinants of fluoroquinolone resistance in a number of Gram-negative pathogens [20], efflux systems of the RND-MFP-OMF type are known to accommodate a broad range of structurally unrelated molecules that can include most classes of antibiotics, as well as biocides, dyes, detergents, metabolic inhibitors, aromatic hydrocarbons (i.e., organic solvents), cationic antimicrobial peptides, toxic fatty acids, bile salts, and homoserine lactones associated with quorum-sensing [13,21]. These last play a role in cell density-dependent expression of a number of virulence factors in (for example) *P. aeruginosa*, and thus the activity of these efflux systems can influence virulence [22,23]. Indeed, a recent study suggests that the MexAB-OprM efflux system of *P. aeruginosa* may promote the release of molecule(s) important for the virulence of this organism [24].

RND-MFP-OMF multidrug efflux systems contributing to antimicrobial resistance have been described in a broad range of Gram-negative pathogens (human, animal, plant) and non-pathogens (Table 1). In view of the role played by RND-MFP-OMF-type multidrug efflux systems in efflux-mediated fluoroquinolone resistance, reports of efflux-mediated resistance to fluoroquinolones in *Vibrio cholerae* [25], *Citrobacter freundii* [26] and *Proteus vulgaris* [27] may also be indicative of multidrug efflux systems in these organisms. The best characterised efflux systems of the RND-MFP-OMF group are those found in *Escherichia coli* [28], *P. aeruginosa* [21] and *Neisseria gonorrhoeae* [29], and homologues of these systems are readily identifiable in both finished genome sequences (see Table 1 for

**Table 1.** RND family multidrug efflux systems of Gram-negative bacteria

Organism	Efflux components <sup>a</sup>			Regulatory genes	Expression <sup>b</sup>	Substrates <sup>c</sup>	Reference(s)
	MFP	RND	OMF				
<i>Acinetobacter baumannii</i>	AdeA	AdeB	AdeC	<i>adeST</i> <sup>d</sup>	wt/+	AG, CX, TC, ER, CM, TP, FQ	[45]
<i>Agrobacterium tumefaciens</i>	AmeA	AmeB	AmeC	<i>ameR</i>	wt/+	BS, NV, SDS	[102]
<i>Brucella suis</i>	AcrA <sup>e</sup> (AAN34087)	AcrB <sup>e</sup> (AAN34086)	?	<i>gntR</i> <sup>f</sup> (AAN34085)	?	?	
<i>Brucella melitensis</i>	AcrA <sup>e</sup> (AAL52073)	AcrB <sup>e</sup> (AAL52074)	?	<i>acrR</i> <sup>f</sup> (AAL52072)	?	?	
<i>Burkholderia cepacia</i>	CeoA (U97042) <sup>g</sup>	CeoB (U97042) <sup>g</sup>	OpcM	?	wt/-; mutant/+	CM, CP, TP	- <sup>h</sup>
<i>Burkholderia pseudomallei</i>	AmrA	AmrB	OprA	<i>amrR</i>	wt/+	ML, AG	- <sup>h</sup>
<i>Campylobacter jejuni</i>	CmeA	CmeB	CmeC	?	wt/+; mutant/?	FQ, ER, BL, RF, TC, EB, SDS, DOC, CM, GN, ACR	[103,104]
<i>Caulobacter crescentus</i>	AcrA <sup>e</sup> (AAK22793)	AcrB <sup>e</sup> (AAK22792)	?	?	?	?	
<i>Enterobacter aerogenes</i>	AcrA	AcrB	TolC	<i>acrR</i>	wt/?; mutant/++	FQ, CM, TC, ACR, NOV, SDS, BS	[32]
<i>Enterobacter cloacae</i>	?	AcrB	?	?		CIP	[33]
<i>Escherichia coli</i> K12	AcrA	AcrB	TolC	<i>acrR, marA, robA, soxS, sdiA</i> <sup>1</sup>	wt/+; <i>marR</i> /++ <sup>j</sup> ; <i>acrR</i> /++	AH <sup>k</sup> , BA, BL, NV, ER, FU, FQ, TC, CM, EB, AC, CV, SDS, TX, BS, TS, PO, FA, MX <sup>k</sup> , LN <sup>k</sup>	- <sup>h</sup>
	AcrE	AcrF	TolC	<i>acrS, sdiA</i> <sup>1</sup>	wt/-; mutant/+	AH, FQ, TC, TP, LN, ML, BS, SDS, AC, RD, BL <sup>m</sup>	- <sup>h</sup>
	AcrA	AcrD	TolC	<i>sdiA</i> <sup>1</sup> , <i>baeR</i> <sup>kk</sup> , <i>cpxR</i> <sup>k</sup> , <i>ompR</i> <sup>kk</sup>	wt/+	AG, NV, DOC, SDS <sup>n</sup> BL <sup>ll</sup>	- <sup>h</sup>
	MdtA (YegM)	MdtB/MdtC (YeN/YegO)	TolC	<i>baeSR</i>	wt/- <sup>o</sup>	NV, BL <sup>ll</sup> , BS, SDS <sup>o</sup>	[54,105,106]
	YhiU	YhiV	TolC	<i>evgA</i>	wt/- <sup>o</sup>	ER, NV, DOX, CV, EB, SDS, BS, BA, RD <sup>o</sup> , BL <sup>ll</sup>	[54,107-109]
<i>Escherichia coli</i> O157:H7	AcrA <sup>P</sup> (BAB33939)	AcrB <sup>P</sup> (BAB33938)	TolC <sup>P</sup> (BAB37346)	<i>acrR</i> <sup>P</sup> (BAB33940)	?	?	
	?	AcrD <sup>P</sup> (BAB36755)	?	?	?	?	
<i>Haemophilus influenzae</i>	AcrA	AcrB	?	?	wt/+	RF, ER, NV, EB, AC, CV, SDS <sup>q</sup>	- <sup>h</sup>
<i>Klebsiella oxytoca</i>	AcrA	AcrB	?	?	wt/?; mutant/+	FQ	[34]
<i>Klebsiella pneumoniae</i>	AcrA <sup>P</sup> (AJ318073)	AcrB <sup>P</sup> (AJ318073)	?	<i>acrR</i> <sup>P</sup> (AJ318073)	wt/?; mutant/+	FQ	[34]
<i>Neisseria gonorrhoeae</i>	MtrC	MtrD	MtrE	<i>mtrR, mtrA</i>	wt/+; <i>mtrR</i> /++ <i>mtrA</i> /-	AZ, ML, RF, PN, TX <sup>r</sup> , CV, CP	- <sup>h</sup>

Table 1. continued

Organism	Efflux components <sup>a</sup>			Regulatory genes	Expression <sup>b</sup>	Substrates <sup>c</sup>	Reference(s)
	MFP	RND	OMF				
<i>Neisseria meningitidis</i>	MtrC (CAB85190)	MtrD (CAB85189)	MtrE (CAB85188)	<i>mtrR</i> (CAB85191), <i>mtrA</i> (CAB85011)	?	?	
<i>Porphyromonas gingivalis</i>	XepA	XepB	XepC	?	wt/+	RF, PM, EB	[110]
<i>Proteus mirabilis</i>	AcrA	AcrB	?	<i>acrR</i>	wt/+	TG, MN, CP, NV, CM, ER, EB, AC, TP, SDS	[80]
<i>Pseudomonas aeruginosa</i>	MexA	MexB	OprM	<i>mexR</i>	wt/+; <i>nalB</i> /+++; <i>nalC</i> <sup>s</sup> /++	BL <sup>t</sup> , FQ, CM, NV, TP, TG <sup>t</sup> , SM, EB, AC, CV, SDS, AH, HL, CL, TL, IR, TS, TPP, RD	— <sup>h</sup>
	MexC	MexD	OprJ	<i>nfxB</i>	wt/— <sup>u</sup> ; <i>nfxB</i> /++	BL <sup>v</sup> , FQ, CM, NV, TP, TG <sup>t</sup> , SM, EB, AC, SDS, AH, CL, TS, TG <sup>v</sup> , TPP, RD	— <sup>h</sup>
	MexE	MexF	OprN	<i>mexT</i>	wt/—; <i>nfxC</i> <sup>w</sup> /++	FQ, CM, TP, AH, TS	— <sup>h</sup>
	MexX (AmrA)	MexY (AmrB)	OprM	<i>mexZ</i> ( <i>amrR</i> )	wt/+	BL <sup>x</sup> , FQ, AG, TC, ER, TG <sup>x</sup>	— <sup>h</sup>
	MexJ	MexK	OprM	<i>mexL</i>	wt/—; mutant/+	TS <sup>y</sup> , ER, TC, CP	[57]
	MexH	MexI	OpmD	?	wt/?; mutant/+	VA, HL?, EB, NOR, RD, AC	[111,111a]
	MexV	MexW	OprM	?	wt/?; mutant/+	FQ, CM, TC, ER, EB, AC	[111b]
	pa1435 <sup>z</sup> (AAG04824)	pa1436 <sup>z</sup> (AAG04825)	?	?	?	?	
	pa3523 <sup>z</sup> (AAG06911)	pa3522 <sup>z</sup> (AAG06910)	pa3521 <sup>z</sup> (AAG06909)	?	?	?	
	pa0156/ pa0157 <sup>z</sup> (AAG03546/ AAG03547)	pa0158 <sup>z</sup> (AAG03548)	?	?	?	?	
	pa2528 <sup>z</sup> (AAG05916)	pa2526/ pa2527 <sup>z</sup> (AAG05914/ AAG05915)	pa2525 <sup>z</sup> (AAG05913)	?	?	?	
	ArpA	ArpB	ArpC	<i>arpR</i>	wt/+	TC, CM, CB, ST, ER, NV	— <sup>h</sup>
	MepA	MepB	MepC	<i>mepR</i>	?	BL, TC, NV, ER, AH	— <sup>h</sup>
<i>Pseudomonas putida</i>	SrpA	SrpB	SrpC	<i>srpR</i> , <i>srpS</i>	wt/+ <sup>aa</sup>	AH	— <sup>h</sup>
	TtgA	TtgB	TtgC	<i>ttgR</i> <sup>ii</sup>	wt/+	CM, AP, TC, TO	— <sup>h</sup>
	TtgD	TtgE	TtgF	?	wt/+ <sup>aa</sup>	AH	— <sup>h</sup>
	TtgG	TtgH	TtgI	<i>ttgV</i> <sup>ij</sup>	wt/+; inducible ++ <sup>bb</sup>	AH, AP, CB	— <sup>h</sup>

Table 1. continued

Organism	Efflux components <sup>a</sup>			Regulatory genes	Expression <sup>b</sup>	Substrates <sup>c</sup>	Reference(s)
	MFP	RND	OMF				
<i>Salmonella enterica</i> serovar Typhimurium	AcrA <sup>P</sup> (AAL19430) ?	AcrB <sup>P</sup> (AAL19429) AcrD <sup>P</sup> (AAL21375)	TolC <sup>P</sup> (AAL22060) ?	<i>acrR</i> <sup>P</sup> (AAL19431) <i>soxRS</i> <sup>cc</sup>	wt/+; mutant/++ <i>soxR</i> /++	BS, SDS, DOC, CH, TX, CV, AC, FA, NV, ER, RF, TC, CM, NOR, NAL, BL, FQ <sup>dd</sup>	— <sup>h</sup>
<i>Salmonella enterica</i> serovar Typhi	AcrA <sup>P</sup> (CAD04961) ?	AcrB <sup>P</sup> (CAD04960) AcrD <sup>P</sup> (CAD07711)	TolC <sup>P</sup> (CAD07712) ?	<i>acrR</i> <sup>P</sup> (CAD04962) ?	? ?		
<i>Serratia marcescens</i>	AcrA	AcrB	?	?	?	FQ	[47,112]
	SdeX	SdeY	?	?	?	ER, TC, NOR, BA, EB, AC, RD	[112a]
<i>Shewanella oneidensis</i>	AcrA <sup>e</sup> (AAN57652)	AcrB <sup>e</sup> (AAN57651)	TolC	?	?	AQDS <sup>ee</sup>	
<i>Shigella flexneri</i>	AcrA <sup>P</sup> (AAN42063)	AcrB <sup>P</sup> (AAN42062)	TolC <sup>P</sup> (AAN44553)	<i>acrR</i> <sup>P</sup> (AAN42064)	?		
<i>Stenotrophomonas maltophilia</i>	SmeA	SmeB	SmeC	<i>smeRS</i>	?	BL, AG, FQ	— <sup>h</sup>
<i>Vibrio cholerae</i>	SmeD ?	SmeE ?	SmeF TolC <sup>hh</sup>	<i>smeT</i> <sup>ff</sup> ?	wt/+ <sup>gg</sup> ; mutant/++ wt/+	TC, ER, FQ, EB, ER, NV, TX	— <sup>h</sup> — <sup>h</sup>
<i>Xanthamonsa campestris</i>	AcrA <sup>e</sup> (AAF96540) AcrA <sup>e</sup> (AAM41645) MexA <sup>e</sup> (AAM41955)	AcrB <sup>e</sup> (AAF96539) AcrB <sup>e</sup> (AAM41646) MexB <sup>e</sup> (AAM41954)	OprM <sup>e</sup> (AAM41953)	<i>acrR/tetR</i> <sup>f</sup> (AAM41644) <i>acrR/tetR</i> <sup>f</sup> (AAM41956)	? ? ?	? ?	
<i>Xylella fastidiosa</i>	AcrA <sup>e</sup> (AAF85183) AcrA <sup>e</sup> (AAF83052)	AcrB <sup>e</sup> (AAF85184) AcrB <sup>e</sup> (AAF83056)	? ?	? ?	? ?	? ?	
<i>Yersinia pestis</i>	AcrA <sup>P</sup> (CAC92367; AAM84631) AcrA <sup>e</sup> (AAM84290)	AcrB <sup>P</sup> (CAC92368; AAM84630) AcrB <sup>e</sup> (AAM84291) AcrD <sup>P</sup> (CAC92285)	TolC <sup>P</sup> (CAC89517) ? ? ?	<i>acrR</i> <sup>P</sup> (CAC92366; AAM84632) ? ? ?	? ? ? ?	? ? ?	

<sup>a</sup>Where the efflux genes/gene products have not been published or their role in antimicrobial efflux and resistance has not been verified, the GenBank protein (three-letter prefix) or gene (one-letter prefix) accession number is shown in parentheses.

<sup>b</sup>wt/+, efflux system is known to be expressed in wild-type cells (under laboratory growth conditions); wt/+ mutant/++, efflux system is expressed in wild-type cells but expression is enhanced in resistant strains; wt/− mutant/+, efflux system is not expressed in wild-type cells but is expressed in resistant strains. In instances where the nature of the mutation leading to enhanced efflux gene expression is known, the gene is indicated along with the relative level of gene expression.

<sup>c</sup>AC, acriflavine; ACR, acridines; AG, aminoglycosides; AH, aromatic hydrocarbons; AP, ampicillin; AQDS, anthraquinone-2,6-disulphonate; AZ, azithromycin; BA, benzalkonium; BL, β-lactams; BS, bile salts; CB, carbenicillin; CH, cholate; CL, cerulenin; CM, chloramphenicol; CP, ciprofloxacin; CV, crystal violet; CX, cefotaxime; DOC, deoxycholate; DOX, doxyrubicin; EB, ethidium bromide; ER, erythromycin; FA, fatty acids; FU, fusaric acid; FQ, fluoroquinolones; GN, gentamicin; HL, homoserine lactones; IR, irgasan; LN, linezolid; ML, macrolides; MN, minocycline; MX, methotrexate; NAL, nalidixic acid; NOR, norfloxacin; NV, novobiocin; PM, puromycin; PN, penicillin; PO, pine oil; RD, rhodamine; RF, rifampicin; SDS, sodium dodecyl sulphate; SM, sulphonamides; ST, streptomycin; TC, tetracycline; TG, tigecycline; TL, thiolactomycin; TO, toluene; TS, triclosan; TP, trimethoprim; TPP, tetraphenyl phosphonium; TX, Triton X-100; VA, vanadium. In instances where only one member of a class of antimicrobial has been tested or is known to be a substrate for a given pump, that member is identified. Where several members of an antimicrobial class are known to be substrates, the class is identified rather than the actual compounds tested.

**Table 1.** footnotes continued

- <sup>d</sup>The *adeST* genes, encoding homologues of the sensor kinase/response regulator members of the superfamily of bacterial two-component regulators, were identified upstream of the *ameABC* efflux genes and implicated in regulating efflux gene expression. No experimental evidence supporting this was provided.
- <sup>e</sup>Homologues of the *E. coli* AcrAB efflux components were identified by a tBlastn search of the genome sequences of the indicated organisms and the corresponding GenBank protein accession number shown in parentheses.
- <sup>f</sup>Examination of gene sequences flanking the corresponding *acrAB*-like efflux genes in the genome sequences of the indicated organisms identified homologues of regulatory genes of the indicated family upstream of the efflux genes, implicating them in the regulation of these efflux genes.
- <sup>g</sup>The *ceoAB* genes (accession number provided in parentheses) and their contribution to efflux-mediated multiresistance genes have not yet been published.
- <sup>h</sup>Except where indicated, accession numbers and references in support of efflux/regulatory gene identification and substrate profiles are provided in recent review articles [13,14,20,21].
- <sup>i</sup>*sdiA* encodes a quorum-sensing regulator that positively regulates efflux gene expression [113,114].
- <sup>j</sup>MarR negatively regulates *marA* gene expression, such that mutations in *marR* (in so-called *mar* mutants) lead to increased MarA production and, thus, increased *acrAB* expression.
- <sup>k</sup>Contribution of AcrAB-TolC to resistance to aromatic hydrocarbons [115], methotrexate [116] and linezolid [117,117a] has recently been confirmed.
- <sup>l</sup>*sdiA* encodes a quorum-sensing regulator that positively influences efflux gene expression.
- <sup>m</sup>Contribution of AcrEF-TolC to resistance to antibiotics [117], dyes and detergents [54] and aromatic hydrocarbons [118] has recently been demonstrated.
- <sup>n</sup>Contribution of AcrD to resistance to non-aminoglycosides has recently been reported [54].
- <sup>o</sup>The efflux genes are not expressed and do not contribute to antimicrobial resistance in wild-type cells, although their cloning on a multicopy plasmid promotes resistance to (and probably efflux of) the indicated agents.
- <sup>p</sup>Orthologues of the *acrAB* efflux genes were identified by a tBlastn search of the genome sequences of the indicated organisms and the corresponding GenBank protein accession number shown in parentheses.
- <sup>q</sup>Data concerning substrate profile were ascertained by examining the susceptibility of an *acrAB* deletion mutant. It is not clear, however, if the apparent differences in substrate profiles between AcrAB in *H. influenzae* and its homologues in *E. coli* and *S. enterica* are real, or whether this pump actually accommodates many more substrates but its highly permeable outer membrane compromises measurable net efflux in this organism.
- <sup>r</sup>Full MtrCDE-mediated resistance to Triton X-100 requires the product of the recently identified *mtrF* gene, encoding a probable cytoplasmic membrane protein of 56 kDa containing 12 transmembrane domains [119].
- <sup>s</sup>The *nalC* gene (also known as pa372) as designated by the *Pseudomonas* Genome Project (<http://www.pseudomonas.com>) is unlinked to the *mexAB-oprM* genes, but encodes a TetR family repressor whose inactivation leads to *mexAB-oprM* hyperexpression.
- <sup>t</sup>MexAB-OprM accommodates many penicillins and cepheims [120], provides modest resistance to a few carbapenems (not imipenem) [121] and is the major efflux system contributing to penem resistance in *P. aeruginosa* [122]. Recently, MexAB-OprM-mediated resistance to the glycylcycline tigecycline has been reported [78].
- <sup>u</sup>Although MexCD-OprJ is not expressed in wild-type cells under normal growth conditions, it is inducible by several of its known substrates, including ethidium bromide, acriflavine, rhodamine 6G, tetraphenylphosphonium chloride, benzalkonium chloride and chlorhexidine [84,84a].
- <sup>v</sup>MexCD-OprJ accommodates many penicillins and cepheims [120], provides modest resistance to a limited number of carbapenems (not imipenem) [121], and promotes modest resistance to penems in the absence of MexAB-OprM [122]. A contribution to tigecycline resistance has also now been reported [78].
- <sup>w</sup>The *nfxC* gene has not yet been identified.
- <sup>x</sup>MexXY-OprM accommodates several penicillins and cepheims [120] and provides modest resistance to a limited number of carbapenems (not imipenem) [121].
- <sup>y</sup>Efflux of triclosan, but not the other antimicrobials, is provided by MexJK alone without OprM involvement.
- <sup>z</sup>Homologues of the known *P. aeruginosa* multidrug efflux systems were identified by a blast search of the *P. aeruginosa* genome sequence and are identified by the 'pa' designations provided by the *Pseudomonas* Genome Project (<http://www.pseudomonas.com>) and the GenBank protein accession numbers (in parentheses).
- <sup>aa</sup>Expression is inducible by aromatic hydrocarbons.
- <sup>bb</sup>Expressed in wild-type cells with enhanced expression occurring in the presence of aromatic hydrocarbons.
- <sup>cc</sup>See Koutsolioutsou *et al.* [123].
- <sup>dd</sup>Contribution of AcrAB-TolC to fluoroquinolone resistance has recently been reported [124].
- <sup>ee</sup>TolC-dependent export of AQDS (anthraquinone-2,6-disulphonate) has been demonstrated [125], although it is not clear whether by a mechanism that involves the AcrAB homologues identified in *Shewanella oneidensis*.
- <sup>ff</sup>*smeT* encodes a repressor of *smeDEF* expression [126].
- <sup>gg</sup>*SmeDEF* contributes to intrinsic multiresistance in *S. maltophilia* [127].
- <sup>hh</sup>While TolC has been implicated in resistance to the indicated agents in *V. cholerae*, it is unclear if the AcrAB homologues identified in this organism work with TolC in providing this resistance.
- <sup>ii</sup>Mediates drug inducibility of TTgABC [127a].
- <sup>jj</sup>Encodes a repressor of *tigGHI* expression [127b].
- <sup>kk</sup>See [127c]. <sup>ll</sup>see [127d].

**Table 2.** Non-RND family multidrug efflux systems of Gram-negative bacteria

Source	Efflux components				Regulatory gene(s)	Expression <sup>a</sup>	Substrates <sup>b</sup>	Reference(s)
	ABC	MATE	MFS	SMR				
Chromosomal								
<i>Bacteroides fragilis</i>	–	–	NorA?	–	?	wt/+; mutant/++	NOR, EB, PM	– <sup>c</sup>
<i>Bacteroides thetaiotaomicron</i>	BexA	–	–	–	?	wt/+; mutant?	FQ, EB	[128]
<i>Bordetella pertussis</i>	–	–	–	BPsmr	?	?	MV, TPP, EB, ACR	[129]
<i>Brucella melitensis</i>	–	NorMI	–	–	?	?	FQ, GN, TPP, ACR, BER	[130]
<i>Burkholderia cepacia</i>	–	–	BcrA	–	?	?	TC, NAL	[131]
<i>Burkholderia vietnamiensis</i>	–	NorM	–	–	?	?	NOR	[132]
<i>Escherichia coli</i>	–	–	EmrB <sup>d</sup>	–	<i>emrR</i> ( <i>mprA</i> )	wt/–; mutant/? <sup>e</sup>	NAL, PMA, HU, TL	– <sup>c</sup>
	–	–	EmrD	–	?	wt/–; mutant/? <sup>e</sup>	HU	– <sup>c</sup>
	–	–	–	EmrE (MvrC)	?	wt/+	EB, ACR, MPP, MV, TC, TPP	– <sup>c</sup>
	–	–	MdfA (CmlA and Cmr)	–	?	wt/+	AG, BA, CM, – <sup>c</sup> DM, EB, ER, FQ, IPTG, PM, RD, RF, TC, TPP	– <sup>c</sup>
	–	–	–	TehA <sup>f</sup>	?	wt/–; mutant/? <sup>e</sup>	CV, EB, PF, TPC	– <sup>c</sup>
	–	YdhE (NorM)	–	–	?	?	NOR, CP, AC, TPP, AG	– <sup>c</sup>
<i>Neisseria gonorrhoeae</i>		NorM			?	?	EB, AC, NOR, CP, BA	[133]
<i>Neisseria meningitidis</i>	–	NorM	–	–	?	?	EB, AC, NOR, CP, BA	[133]
<i>Pseudomonas aeruginosa</i>	–	–	–	EmrE	?	wt/+	EB, ACR, MV, AG	[129,134]
<i>Vibrio cholerae</i>	–	–	VceB <sup>g</sup>	–	?	wt/+	DOC, CCCP, CM, ER, PMA, PCP, NAL	– <sup>c</sup>
<i>Vibrio parahaemolyticus</i>	–	NorM <sup>h</sup>	–	–	?	?	NOR, CP, AG, EB	– <sup>c</sup>
		VmrA			?	?	TPP, ACR, EB	[135]
Plasmid-encoded								
	–	–	–	QacE	?	NA <sup>j</sup>	EB, PF, CV, QAC	– <sup>c</sup>
	–	–	–	QacEA1	?	NA <sup>j</sup>	EB, PF, CV, QAC	– <sup>c</sup>
	McbF <sup>i</sup>	–	–	–	<i>emrR</i> ( <i>mprA</i> )	NA <sup>j</sup>	FQ, HU	– <sup>c</sup>

<sup>a</sup>Relative expression of the corresponding efflux system is indicated in wild-type (wt) cells (where known) and in mutants known to express the system. In instances where the nature of the mutation leading to enhanced efflux gene expression is known, the gene is indicated along with the relative level of gene expression. wt/+ mutant/++, efflux system is expressed in wild-type cells (under laboratory growth conditions) but expression is enhanced in resistant strains; wt/– mutant/+, efflux system is not expressed in wild-type cells but is expressed in resistant strains; ?, expression status unknown.

Table 2. footnotes continued

<sup>b</sup>AC, acriflavine; ACR, acridines; AG, aminoglycosides; BA, benzalkonium; BER, berberine; CCCP, carbonyl cyanide *m*-chlorophenylhydrazone; CM, chloramphenicol; CP, ciprofloxacin; CV, crystal violet; DM, daunomycin; DOC, deoxycholate; EB, ethidium bromide; ER, erythromycin; FQ, fluoroquinolones; GN, gentamicin; HU, hydrophobic uncouplers; IPTG, isopropyl- $\beta$ -D-thiogalactopyranoside; MPP, methyltriphenylphosphonium; MV, methyl viologen; NAL, nalidixic acid; NOR, norfloxacin; PCP, pentachlorophenol; PF, proflavin; PM, puromycin; PMA, phenylmercuric acetate; QAC, quaternary ammonium compounds; RD, rhodamine; RF, rifampicin; TC, tetracycline; TL, thiolactomycin; TPC, tetraphenylarsonium chloride; TPP, tetraphenyl phosphonium. In instances where only one member of a class of antimicrobial has been tested or is known to be a substrate for a given pump, that member is identified. Where several members of an antimicrobial class are known to be substrates, the class is identified rather than the actual compounds tested.

<sup>c</sup>Accession numbers and references in support of efflux/regulatory gene identification and substrate profiles are provided in recent review articles [13,14,20].

<sup>d</sup>Part of a probable three-component export system that includes the MFP component, EmrA, and the OMF component, TolC.

<sup>e</sup>Efflux activity and/or drug resistance has only been shown for the gene cloned on a plasmid.

<sup>f</sup>Part of a two-component export system that includes TehB.

<sup>g</sup>Part of a two-component export system that includes the MFP component VceA.

<sup>h</sup>First described in *V. parahaemolyticus*, NorM homologues (ascertained using BLAST searches) are found in the genome sequences of a broad range of Gram-negative organisms, including most of those listed in Table 1.

<sup>i</sup>Part of a two-component export system that includes McbE.

<sup>j</sup>Not applicable.

examples) and unfinished genome sequences (e.g., *Bordetella pertussis*, *Bordetella parapertussis*, *Bordetella bronchiseptica*, *Burkholderia mallei*, *Legionella pneumophila*, *Pasteurella multocida* and *Pseudomonas syringae*; [http://www.ncbi.nlm.nih.gov/cgi-bin/Entrez/genom\\_table.cgi](http://www.ncbi.nlm.nih.gov/cgi-bin/Entrez/genom_table.cgi)) of a number of Gram-negative bacteria, although their contribution to antimicrobial resistance in these organisms remains to be determined. Increasingly, multidrug efflux systems of the RND-MFP-OMF type are implicated in clinical episodes involving antibiotic-resistant organisms. Examples include *E. coli* [30,31], *Enterobacter aerogenes* [32,32a], *Enterobacter cloacae* [33], *Klebsiella* spp. [34], *N. gonorrhoeae* [35–37], *Salmonella enterica* serovar Typhimurium [38–41,41a], *P. aeruginosa* [42,43,43a,b], *S. maltophilia* [44], *Acinetobacter baumannii* [45], *Burkholderia cepacia* [46] and, possibly, *Serratia marcescens* [47] and *Campylobacter jejuni* [48]. Additional references can be found in other recent review articles [13,20,21]. Efflux-mediated fluoroquinolone resistance in clinical isolates of *C. freundii* [26], *Ent. aerogenes* [49], *Ent. cloacae* [33] and *Klebsiella pneumoniae* [50,51] may also be attributable to multidrug efflux systems.

## ANTIBIOTIC-BIocide CROSS-RESISTANCE

The observation that RND-MFP-OMF efflux systems in both *E. coli* (AcrAB [52–54]) and *P. aeruginosa* (MexAB-OprM [55], MexCD-OprJ

[56], MexEF-OprN [56] and MexJK [57]) accommodate both biocides and clinically important antibiotics has added fuel to the debate on biocide-antibiotic co-resistance and the possible selection of multiply antibiotic-resistant strains by biocides [9,58–64,64a]. While it appears that such multidrug efflux systems are unlikely to facilitate resistance to biocides used at the recommended ‘working’ concentrations of these agents, the risk of residual biocide levels selecting pump-producing mutants that are concomitantly resistant to clinically relevant antibiotics is of some concern [65]. Certainly, in-vitro studies have confirmed that biocide-resistant strains which overexpress multidrug efflux systems, and thus become resistant to multiple antibiotics, can be selected in both *E. coli* [52,53] and *P. aeruginosa* [56]. Given the broad distribution of Mex- and Acr-like efflux systems in Gram-negative bacteria (see above and Table 1), the risk of efflux-mediated biocide-antibiotic cross-resistance in other pathogens is a real possibility.

## EFFLUX SYNERGY WITH OTHER RESISTANCE MECHANISMS

The intrinsic and acquired resistance of Gram-negative bacteria invariably depends upon reduced antimicrobial accumulation, with or without antimicrobial modification or destruction or target site alterations [13]. The former comes about as a result of the synergy between the



outer-membrane barrier and active efflux mechanisms, and disruption of either is effective in enhancing accumulation and, thus, susceptibility [66]. The importance of this synergy is highlighted by the concomitant decline in porin production, which decreases outer-membrane permeability to, for example, hydrophilic agents, and the increase in efflux activity that is seen in so-called multiple antibiotic-resistant or *mar* mutants of *E. coli* [67]. A *mar* locus (or a Mar-like phenotype) is, in fact, widely distributed among Gram-negative bacteria [13,68,69]. Increasingly, resistance to fluoroquinolones in Gram-negative bacteria results from a combination of enhanced expression of multidrug efflux systems and target site mutations [25,39,48,70–73]. It may be that efflux was also a contributing (but overlooked) factor in earlier reports of fluoroquinolone resistance attributed to target site mutations. Significantly, loss of multidrug efflux (AcrAB in *E. coli* [74] or *Salmonella typhimurium* [39]; MexAB–OprM in *P. aeruginosa* [75]; CmeABC in *Camp. jejuni*) can undermine the resistance provided by target site mutations, in some instances rendering the mutants susceptible despite the presence of these target site mutations [39,74,75]. These results highlight both the significance of the contribution of efflux mechanisms to fluoroquinolone resistance in Gram-negative organisms, and the probable utility of efflux inhibitors in countering resistance attributable to efflux and target site mutations.

## OVERCOMING EFFLUX-MEDIATED MULTIRESISTANCE

An obvious approach to overcoming resistance mediated by efflux mechanisms is to target them directly (i.e., develop efflux inhibitors) or to develop antimicrobials that are less impacted by efflux (i.e., are poor substrates for these efflux systems). To date, several inhibitors active against the multidrug efflux systems of Gram-negative bacteria have been described that are, for example, effective at reversing efflux-mediated multiresistance [4,76,77]. The glycylicycline tigecycline (GAR-936), originally identified as being unaffected by (for example) the tetracycline-specific Tet efflux determinants found in *E. coli*, also appears to be an inferior substrate for the multidrug efflux systems of *P. aeruginosa* [78] and is active against multiresistant *S. maltophilia* and *A. baumannii* [79].

Nonetheless, *P. aeruginosa* is still generally resistant to this agent, which is exported by the MexAB–OprM, MexCD–OprJ and MexXY–OprM multidrug efflux systems of this organism [78]. Moreover, the notably reduced susceptibility of *Proteus mirabilis* to tigecycline was recently explained by the activity of the AcrAB multidrug efflux system of this organism, which readily accommodates this agent [80]. The likelihood that many novel agents may be substrates for these highly accommodating efflux systems [81] needs to inform current antimicrobial strategies that focus on developing novel agents or novel drug targets [82]. Given the synergy that exists between outer-membrane impermeability and active drug efflux, targeting the outer-membrane barrier is also proposed as an effective way of compromising efflux-mediated multiresistance [13]. Finally, therapeutic approaches based on vaccines, immunomodulation, cationic antibacterial peptides and phages, to name just a few, may become increasingly important in treating multiresistant Gram-negative infections, benefiting as they do from an ‘insensitivity’ to drug efflux mechanisms [4].

## NATURAL FUNCTION OF MULTIDRUG EFFLUX SYSTEMS

There is much discussion in the literature as to the natural function of bacterial multidrug efflux systems, with evidence for induction of the systems by agents known to be exported, providing support for proposed roles in protection against noxious exogenous substances. For example, the MexXY–OprM system of *P. aeruginosa* is both induced by and exports several antibiotics, including gentamicin, erythromycin and tetracycline [83], and, as such, efflux of these agents may be its intended function. However, it is also possible that the action of these agents on their ribosome targets stimulates the production of cellular product(s) which are the intended MexXY–OprM substrates and, thus, induce expression of this efflux. The MexCD–OprJ efflux system of this same organism is also inducible by a number of its non-antibiotic substrates (e.g., the dyes rhodamine 6G, acriflavine and ethidium bromide and the biocides benzalkonium chloride and chlorhexid), although not by any clinically relevant antibiotics [84a]. The observed inducibility of the *E. coli* AcrAB system by toxic fatty acids [85], and the demonstrated role of AcrAB in the export of and

resistance to bile salts [86], have been cited as support for a role for AcrAB in protecting the cell from the action of these agents in the gut [85], although clearly there is no evidence for antibiotics being the intended substrate. Similarly, a protective function has been attributed to the MtrCDE system, which provides resistance to faecal lipids in rectal isolates of *N. gonorrhoeae* [87] and, probably, bile salts known to bathe mucous membranes [88]. Still, the ability to accommodate bile salts is not, in itself, proof of a protective function for either efflux system, inasmuch as many of the so-called multidrug efflux systems of (for example) *E. coli* accommodate and provide resistance to bile salts [54]. It would appear unlikely that a given organism would devote several systems to the export of bile salts, and so these agents may be just some of many, perhaps unintended, substrates for these highly accommodating, broadly specific efflux systems.

Export of bacterial products is also a proposed function for these multidrug efflux systems, with a previous review implicating byproducts of metabolism as probable substrates for these efflux systems [89]. The upregulation of the *E. coli* AcrAB efflux system in strains with mutations in central biosynthetic pathways, possibly as a result of accumulation of pathway intermediates [90], certainly support this idea. The fact that a homologue, IefABC, of the AcrAB and MexAB efflux systems exports isoflavanoids in the plant pathogen *Agrobacterium tumefaciens* adds support to suggestions that non-antimicrobials are the intended substrates of many of these efflux systems [91]. Finally, efflux homologues of the RND-MFP-OMF type are broadly conserved among Gram-negative organisms, with multiple examples present in many species (Table 1) that are invariably linked to regulatory genes, implying individual control of these systems and, thus, distinct functions in the cell. So, while these systems, even in a single organism, accommodate many of the same antimicrobials, it is unlikely that their export is the primary function of these systems. Still, given their contribution to resistance against many clinically relevant agents, and their broad distribution, it is clear that these efflux systems are important determinants of antimicrobial resistance in many Gram-negative pathogens, and that their operation in these organisms is likely to compromise the use of current and yet to be developed agents.

## CONCLUSIONS

Multiresistance is a significant problem in medicine because it limits chemotherapeutic options. The recognition that many instances of intrinsic and acquired antimicrobial (including biocide) resistance are attributable, at least in part, to the activity of broadly specific multidrug efflux systems, and that these are broadly conserved in Gram-negative pathogens, means that antimicrobial (including biocide) use and development must be informed by the possible impact of efflux. While multidrug efflux might compromise the activity of novel agents that could be efflux substrates, a possible bigger concern is that potent new agents might be missed in, for example, whole cell screens precisely because they are excluded from the cell by these efflux mechanisms [81]. Current efforts aimed at developing multidrug efflux inhibitors [77,92] that can be used in combination with existing and, possibly, novel agents (a strategy that has proven effective in the case of  $\beta$ -lactam- $\beta$ -lactamase inhibitors [93]) may ultimately prove important in preserving a strategy for treating infectious disease based on chemotherapeutic agents. Such efforts will be aided by recent reports of the three-dimensional structures of components of the RND-MFP-OMF pump family [94–96], and by the identification of the drug recognition regions of the RND component [97–101], both of which can assist in rational inhibitor design.

## REFERENCES

1. Waterer GW, Wunderink RG. Increasing threat of Gram-negative bacteria. *Crit Care Med* 2001; **29**: N75–N81.
2. Jones RN. Resistance patterns among nosocomial pathogens: trends over the past few years. *Chest* 2001; **119**: 397S–404S.
3. Livermore DM. Bacterial resistance: origins, epidemiology, and impact. *Clin Infect Dis* 2003; **36**: S11–S23.
4. Poole K. Overcoming multidrug resistance in Gram-negative bacteria. *Curr Opin Invest Drugs* 2003; **4**: 128–139.
5. Hosein IK, Hill DW, Jenkins LE, Magee JT. Clinical significance of the emergence of bacterial resistance in the hospital environment. *J Appl Microbiol* 2002; **92**(suppl 1): 90S–97S.
6. Helms M, Vastrup P, Gerner-Smidt P, Molbak K. Excess mortality associated with antimicrobial drug-resistant *Salmonella typhimurium*. *Emerg Infect Dis* 2002; **8**: 490–495.
7. Carmeli Y, Troillet N, Karchmer AW, Samore MH. Health and economic outcomes of antibiotic resistance in

- Pseudomonas aeruginosa*. *Arch Intern Med* 1999; **159**: 1127–1132.
8. Niederman MS. Impact of antibiotic resistance on clinical outcomes and the cost of care. *Crit Care Med* 2001; **29**: N114–N120.
9. Poole K. Mechanisms of bacterial biocide and antibiotic resistance. *J Appl Microbiol* 2002; **92**(suppl 1): 55S–64S.
10. Hogan D, Kolter R. Why are bacteria refractory to antimicrobials? *Curr Opin Microbiol* 2002; **5**: 472–477.
11. Normark BH, Normark S. Evolution and spread of antibiotic resistance. *J Intern Med* 2002; **252**: 91–106.
12. Rowe-Magnus DA, Davies J, Mazel D. Impact of integrons and transposons on the evolution of resistance and virulence. *Curr Top Microbiol Immunol* 2002; **264**: 167–188.
13. Poole K. Outer membranes and efflux: the path to multidrug resistance in gram-negative bacteria. *Curr Pharm Biotechnol* 2002; **3**: 77–98.
14. Poole K. Multidrug resistance in Gram-negative bacteria. *Curr Opin Microbiol* 2001; **4**: 500–508.
15. Webber MA, Piddock LJ. The importance of efflux pumps in bacterial antibiotic resistance. *J Antimicrob Chemother* 2003; **51**: 9–11.
16. Paulsen IT, Chen J, Nelson KE, Saier MH. Comparative genomics of microbial drug efflux systems. *J Mol Microbiol Biotechnol* 2001; **3**: 145–150.
17. Putman M, van Veen HW, Konings WN. Molecular properties of bacterial multidrug transporters. *Microbiol Mol Biol Rev* 2000; **64**: 672–693.
18. Zgurskaya HI, Nikaido H. Multidrug resistance mechanisms: drug efflux across two membranes. *Mol Microbiol* 2000; **37**: 219–225.
19. Kobayashi N, Nishino K, Yamaguchi A. Novel macro-lide-specific ABC-type efflux transporter in *Escherichia coli*. *J Bacteriol* 2001; **183**: 5639–5644.
20. Poole K. Efflux-mediated resistance to fluoroquinolones in Gram-negative bacteria. *Antimicrob Agents Chemother* 2000; **44**: 2233–2241.
21. Poole K. Multidrug efflux pumps and antimicrobial resistance in *Pseudomonas aeruginosa* and related organisms. *J Mol Microbiol Biotechnol* 2001; **3**: 255–264.
22. Sanchez P, Linares JF, Ruiz-Diez B *et al*. Fitness of *in vitro* selected *Pseudomonas aeruginosa* *nalB* and *nfxB* multidrug resistant mutants. *J Antimicrob Chemother* 2002; **50**: 657–664.
23. Cosson P, Zulianello L, Join-Lambert O *et al*. *Pseudomonas aeruginosa* virulence analyzed in a *Dictyostelium discoideum* host system. *J Bacteriol* 2002; **184**: 3027–3033.
24. Hirakata Y, Srikumar R, Poole K *et al*. Multidrug efflux systems play an important role in the invasiveness of *Pseudomonas aeruginosa*. *J Exp Med* 2002; **196**: 109–118.
25. Baranwal S, Dey K, Ramamurthy T, Nair GB, Kundu M. Role of active efflux in association with target gene mutations in fluoroquinolone resistance in clinical isolates of *Vibrio cholerae*. *Antimicrob Agents Chemother* 2002; **46**: 2676–2678.
26. Navia MM, Ruiz J, Ribera A, de Anta MT, Vila J. Analysis of the mechanisms of quinolone resistance in clinical isolates of *Citrobacter freundii*. *J Antimicrob Chemother* 1999; **44**: 743–748.
27. Ishii H, Sato K, Hoshino K *et al*. Active efflux of ofloxacin by a highly quinolone-resistant strain of *Proteus vulgaris*. *J Antimicrob Chemother* 1991; **28**: 827–836.
28. Nikaido H, Zgurskaya HI. AcrAB and related multidrug efflux pumps of *Escherichia coli*. *J Mol Microbiol Biotechnol* 2001; **3**: 215–218.
29. Shafer WM, Veal WL, Lee EH, Zarantonelli L, Balthazar JT, Rouquette C. Genetic organization and regulation of antimicrobial efflux systems possessed by *Neisseria gonorrhoeae* and *Neisseria meningitidis*. *J Mol Microbiol Biotechnol* 2001; **3**: 219–224.
30. Mazzariol A, Tokue Y, Kanegawa TM, Cornaglia G, Nikaido H. High-level fluoroquinolone-resistant clinical isolates of *Escherichia coli* overproduce multidrug efflux protein AcrA. *Antimicrob Agents Chemother* 2000; **44**: 3441–3443.
31. Linde HJ, Notka F, Metz M, Kochanowski B, Heisig P, Lehn N. In vivo increase in resistance to ciprofloxacin in *Escherichia coli* associated with deletion of the C-terminal part of MarR. *Antimicrob Agents Chemother* 2000; **44**: 1865–1868.
32. Pradel E, Pages JM. The AcrAB–TolC efflux pump contributes to multidrug resistance in the nosocomial pathogen *Enterobacter aerogenes*. *Antimicrob Agents Chemother* 2002; **46**: 2640–2643.
- 32a. Gayet S, Chollet R, Molle G, Pages JM, Chevalier J. Modification of outer membrane protein profile and evidence suggesting an active drug pump in *Enterobacter aerogenes* clinical strains. *Antimicrob Agents Chemother* 2003; **47**: 1555–1559.
33. Linde HJ, Notka F, Irtenkauf C *et al*. Increase in MICs of ciprofloxacin in vivo in two closely related clinical isolates of *Enterobacter cloacae*. *J Antimicrob Chemother* 2002; **49**: 625–630.
34. Mazzariol A, Zuliani J, Cornaglia G, Rossolini GM, Fontana R. AcrAB efflux system: expression and contribution to fluoroquinolone resistance in *Klebsiella* spp. *Antimicrob Agents Chemother* 2002; **46**: 3984–3986.
35. Zarantonelli L, Borthagaray G, Lee EH, Veal W, Shafer WM. Decreased susceptibility to azithromycin and erythromycin mediated by a novel *mtr* (R) promoter mutation in *Neisseria gonorrhoeae*. *J Antimicrob Chemother* 2001; **47**: 651–654.
36. Veal WL, Nicholas RA, Shafer WM. Overexpression of the MtrC–MtrD–MtrE efflux pump due to an *mtrR* mutation is required for chromosomally mediated penicillin resistance in *Neisseria gonorrhoeae*. *J Bacteriol* 2002; **184**: 5619–5624.
37. Cousin JS, Whittington WL, Roberts MC. Acquired macrolide resistance genes and the 1 bp deletion in the *mtrR* promoter in *Neisseria gonorrhoeae*. *J Antimicrob Chemother* 2003; **51**: 131–133.
38. Randall LP, Cooles SW, Sayers AR, Woodward MJ. Association between cyclohexane resistance in *Salmonella* of different serovars and increased resistance to multiple antibiotics, disinfectants and dyes. *J Med Microbiol* 2001; **50**: 919–924.
39. Baucheron S, Imberechts H, Chaslus-Dancla E, Cloeckert A. The AcrB multidrug transporter plays a major role in high-level fluoroquinolone resistance in *Salmonella enterica* serovar typhimurium phage type DT204. *Microb Drug Resist* 2002; **8**: 281–289.
40. Piddock LJ. Fluoroquinolone resistance in *Salmonella* serovars isolated from humans and food animals. *FEMS Microbiol Rev* 2002; **26**: 3–16.

41. Piddock LJ, White DG, Gensberg K, Pumbwe L, Griggs DJ. Evidence for an efflux pump mediating multiple antibiotic resistance in *Salmonella enterica* serovar Typhimurium. *Antimicrob Agents Chemother* 2000; **44**: 3118–3121.
- 41a. Soto SM, Ruiz J, Mendoza MC, Vila J. In vitro fluoroquinolone-resistant mutants of *Salmonella enterica* serotype Enteritidis: analysis of mechanisms involved in resistance. *Int J Antimicrob Agents* 2003; **22**: 537–540.
42. Jalal S, Ciofu O, Hoiby N, Gotoh N, Wretling B. Molecular mechanisms of fluoroquinolone resistance in *Pseudomonas aeruginosa* isolates from cystic fibrosis. *Antimicrob Agents Chemother* 2000; **44**: 710–712.
43. Cavallo JD, Plesiat P, Couetdic G, Leblanc F, Fabre R. Mechanisms of  $\beta$ -lactam resistance in *Pseudomonas aeruginosa*: prevalence of OprM-overproducing strains in a French multicentre study (1997). *J Antimicrob Chemother* 2002; **50**: 1039–1043.
- 43a. Higgins PG, Fluit AC, Milatovic D, Verhoef J, Schmitz FJ. Mutations in GyrA, ParC, MexR and NfxB in clinical isolates of *Pseudomonas aeruginosa*. *Int J Antimicrob Agents* 2003; **21**: 409–413.
- 43b. Sobel ML, McKay GA, Poole K. Contribution of the MexXY multidrug transporter to aminoglycoside resistance in *Pseudomonas aeruginosa* clinical isolates. *Antimicrob Agents Chemother* 2003; **47**: 3202–3207.
44. Zhang L, Li X-Z, Poole K. Multiple antibiotic resistance in *Stenotrophomonas maltophilia*: involvement of a multidrug efflux system. *Antimicrob Agents Chemother* 2000; **44**: 287–293.
45. Magnet S, Courvalin P, Lambert T. Resistance-nodulation-cell division-type efflux pump involved in aminoglycoside resistance in *Acinetobacter baumannii* strain BM4454. *Antimicrob Agents Chemother* 2001; **45**: 3375–3380.
46. Burns JL, Hedin LA, Lien DM. Chloramphenicol resistance in *Pseudomonas cepacia* because of decreased permeability. *Antimicrob Agents Chemother* 1989; **33**: 136–141.
47. Kumar A, Worobec EA. Fluoroquinolone resistance of *Serratia marcescens*: involvement of a proton gradient-dependent efflux pump. *J Antimicrob Chemother* 2002; **50**: 593–596.
48. Luo N, Sahin O, Lin J, Michel LO, Zhang Q. In vivo selection of *Campylobacter* isolates with high levels of fluoroquinolone resistance associated with *gyrA* mutations and the function of the CmeABC efflux pump. *Antimicrob Agents Chemother* 2003; **47**: 390–394.
49. Mallea M, Chevalier J, Bornet C *et al.* Porin alteration and active efflux: two *in vivo* drug resistance strategies used by *Enterobacter aerogenes*. *Microbiology* 1998; **144**: 3003–3009.
50. Deguchi T, Kawamura T, Yasuda M *et al.* In vivo selection of *Klebsiella pneumoniae* strains with enhanced quinolone resistance during fluoroquinolone treatment of urinary tract infections. *Antimicrob Agents Chemother* 1997; **41**: 1609–1611.
51. Martínez-Martínez L, García I, Ballesta S, Benedí VJ, Hernández-Allés S, Pascual A. Energy-dependent accumulation of fluoroquinolones in quinolone-resistant *Klebsiella pneumoniae* strains. *Antimicrob Agents Chemother* 1998; **42**: 1850–1852.
52. McMurry LM, Oethinger M, Levy SB. Overexpression of *marA*, *soxS*, or *acrAB* produces resistance to triclosan in laboratory and clinical strains of *Escherichia coli*. *FEMS Microbiol Lett* 1998; **166**: 305–309.
53. Moken MC, McMurry LM, Levy SB. Selection of multiple-antibiotic-resistant (Mar) mutants of *Escherichia coli* by using the disinfectant pine oil: roles of the *mar* and *acrAB* loci. *Antimicrob Agents Chemother* 1997; **41**: 2770–2772.
54. Nishino K, Yamaguchi A. Analysis of a complete library of putative drug transporter genes in *Escherichia coli*. *J Bacteriol* 2001; **183**: 5803–5812.
55. Schweizer HP. Intrinsic resistance to inhibitors of fatty acid biosynthesis in *Pseudomonas aeruginosa* is due to efflux: application of a novel technique for generation of unmarked chromosomal mutations for the study of efflux systems. *Antimicrob Agents Chemother* 1998; **42**: 394–398.
56. Chuanchuen R, Beinlich K, Hoang TT, Becher A, Karkhoff-Schweizer RR, Schweizer HP. Cross-resistance between triclosan and antibiotics in *Pseudomonas aeruginosa* is mediated by multidrug efflux pumps: exposure of a susceptible mutant strain to triclosan selects *nfxB* mutants overexpressing MexCD–OprJ. *Antimicrob Agents Chemother* 2001; **45**: 428–432.
57. Chuanchuen R, Narasaki CT, Schweizer HP. The MexJK efflux pump of *Pseudomonas aeruginosa* requires OprM for antibiotic efflux but not for efflux of triclosan. *J Bacteriol* 2002; **184**: 5036–5044.
58. McBain AJ, Rickard AH, Gilbert P. Possible implications of biocide accumulation in the environment on the prevalence of bacterial antibiotic resistance. *J Ind Microbiol Biotechnol* 2002; **29**: 326–330.
59. Fraise AP. Reply. *J Antimicrob Chemother* 2002; **50**: 139–140.
60. Gilbert P, McBain AJ, Bloomfield SF. Biocide abuse and antimicrobial resistance: being clear about the issues. *J Antimicrob Chemother* 2002; **50**: 137–139.
61. Fraise AP. Biocide abuse and antimicrobial resistance—a cause for concern? *J Antimicrob Chemother* 2002; **49**: 11–12.
62. Schweizer HP. Triclosan: a widely used biocide and its link to antibiotics. *FEMS Microbiol Lett* 2001; **202**: 1–7.
63. Ng ME, Jones S, Leong SH, Russell AD. Biocides and antibiotics with apparently similar actions on bacteria: is there the potential for cross-resistance? *J Hosp Infect* 2002; **51**: 147–149.
64. Russell AD. Introduction of biocides into clinical practice and the impact on antibiotic-resistant bacteria. *J Appl Microbiol* 2002; **92**(suppl 1): 121S–135S.
- 64a. Gilbert P, McBain AJ. Potential impact of increased use of biocides in consumer products on prevalence of antibiotic resistance. *Clin Microbiol Rev* 2003; **16**: 189–208.
65. Russell AD. Biocides and pharmacologically active drugs as residues and in the environment: is there a correlation with antibiotic resistance? *Am J Infect Control* 2002; **30**: 495–498.
66. Li X-Z, Zhang L, Poole K. Interplay between the MexAB–OprM multidrug efflux system and the outer membrane barrier in the multiple antibiotic resistance of *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 2000; **45**: 433–436.
67. Randall LP, Woodward MJ. The multiple antibiotic resistance (*mar*) locus and its significance. *Res Vet Sci* 2002; **72**: 87–93.

68. Cohen SH, Yan W, Levy SB. A multidrug resistance regulatory chromosomal locus is widespread among enteric bacteria. *J Infect Dis* 1993; **168**: 484–488.
69. Chollet R, Bollet C, Chevalier J, Mallea M, Pages JM, Davin-Regli A. *mar* operon involved in multidrug resistance of *Enterobacter aerogenes*. *Antimicrob Agents Chemother* 2002; **46**: 1093–1097.
70. Nakajima A, Sugimoto Y, Yoneyama H, Nakae T. High-level fluoroquinolone resistance in *Pseudomonas aeruginosa* due to interplay of the MexAB–OprM efflux pump and the DNA gyrase mutation. *Microbiol Immunol* 2002; **46**: 391–395.
71. Tavio M, Vila J, Ruiz J *et al.* In vitro selected fluoroquinolone-resistant mutants of *Citrobacter freundii*: analysis of the quinolone resistance acquisition. *J Antimicrob Chemother* 2000; **45**: 521–524.
72. Le TI, Couetdic G, Clermont O, Brahimi N, Plesiat P, Bingen E. In vivo selection of a target/efflux double mutant of *Pseudomonas aeruginosa* by ciprofloxacin therapy. *J Antimicrob Chemother* 2001; **48**: 553–555.
73. Martinez-Martinez L, Pascual A, Conejo MC *et al.* Energy-dependent accumulation of norfloxacin and porin expression in clinical isolates of *Klebsiella pneumoniae* and relationship to extended-spectrum  $\beta$ -lactamase production. *Antimicrob Agents Chemother* 2002; **46**: 3926–3932.
74. Oethinger M, Kern WV, Jellen-Ritter AS, McMurphy LM, Levy SB. Ineffectiveness of topoisomerase mutations in mediating clinically significant fluoroquinolone resistance in *Escherichia coli* in the absence of the AcrAB efflux pump. *Antimicrob Agents Chemother* 2000; **44**: 10–13.
75. Lomovskaya O, Lee A, Hoshino K *et al.* Use of a genetic approach to evaluate the consequences of inhibition of efflux pumps in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 1999; **43**: 1340–1346.
76. Poole K. Overcoming antimicrobial resistance by targeting resistance mechanisms. *J Pharm Pharmacol* 2001; **53**: 283–294.
77. Lomovskaya O, Watkins W. Inhibition of efflux pumps as a novel approach to combat drug resistance in bacteria. *J Mol Microbiol Biotechnol* 2001; **3**: 225–236.
78. Dean CR, Visalli MA, Projan SJ, Sum P-E, Bradford PA. Efflux-mediated resistance to tigecycline (GAR-936) in *Pseudomonas aeruginosa* PAO1. *Antimicrob Agents Chemother* 2003; **47**: 972–978.
79. Betriu C, Rodriguez-Avial I, Sanchez BA, Gomez M, Alvarez J, Picazo JJ. In vitro activities of tigecycline (GAR-936) against recently isolated clinical bacteria in Spain. *Antimicrob Agents Chemother* 2002; **46**: 892–895.
80. Visalli MA, Murphy E, Projan SJ, Bradford PA. AcrAB multidrug efflux pump is associated with reduced levels of susceptibility to tigecycline (GAR-936) in *Proteus mirabilis*. *Antimicrob Agents Chemother* 2003; **47**: 665–669.
81. Tegos G, Stermitz FR, Lomovskaya O, Lewis K. Multidrug pump inhibitors uncover remarkable activity of plant antimicrobials. *Antimicrob Agents Chemother* 2002; **46**: 3133–3141.
82. Ali ST, Gill AE, Lewendon A. Novel targets for antibiotic drug design. *Curr Opin Invest Drugs* 2002; **3**: 1712–1717.
83. Masuda N, Sakagawa E, Ohya S, Gotoh N, Tsujimoto H, Nishino T. Contribution of the MexX–MexY–OprM efflux system to intrinsic resistance in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2000; **44**: 2242–2246.
84. Morita Y, Komori Y, Mima T, Kuroda T, Mizushima T, Tsuchiya T. Construction of a series of mutants lacking all of the four major *mex* operons for multidrug efflux pumps or possessing each one of the operons from *Pseudomonas aeruginosa* PAO1: MexCD–OprJ is an inducible pump. *FEMS Microbiol Lett* 2001; **202**: 139–143.
- 84a. Morita Y, Murata T, Mima T *et al.* Induction of *mexCD-oprJ* operon for a multidrug efflux pump by disinfectants in wild-type *Pseudomonas aeruginosa* PAO1. *J Antimicrob Chemother* 2003; **5**: 991–994.
85. Ma D, Cook DN, Alberti M, Pon NG, Nikaido H, Hearst JE. Genes *acrA* and *acrB* encode a stress-induced system of *Escherichia coli*. *Mol Microbiol* 1995; **16**: 45–55.
86. Thanassi DG, Cheng LW, Nikaido H. Active efflux of bile salts in *Escherichia coli*. *J Bacteriol* 1997; **179**: 2512–2518.
87. Shafer WM, Balthazar JT, Hagman KE, Morse SA. Missense mutations that alter the DNA-binding domain of the MtrR protein occur frequently in rectal isolates of *Neisseria gonorrhoeae* that are resistant to fecal lipids. *Microbiology* 1995; **141**: 907–911.
88. Hagman KE, Pan W, Spratt BG, Balthazar JT, Judd RC, Shafer WM. Resistance of *Neisseria gonorrhoeae* to antimicrobial hydrophobic agents is modulated by the *mtrRCDE* efflux system. *Microbiology* 1995; **141**: 611–622.
89. Poole K. Bacterial multidrug resistance—emphasis on efflux mechanisms and *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 1994; **34**: 453–456.
90. Helling RB, Janes BK, Kimball H *et al.* Toxic waste disposal in *Escherichia coli*. *J Bacteriol* 2002; **184**: 3699–3703.
91. Palumbo JD, Kado CI, Phillips DA. An isoflavonoid-inducible efflux pump in *Agrobacterium tumefaciens* is involved in competitive colonization of roots. *J Bacteriol* 1998; **180**: 3107–3113.
92. Ryan BM, Dougherty TJ, Beaulieu D, Chuang J, Dougherty BA, Barrett JF. Efflux in bacteria: what do we really know about it? *Expert Opin Invest Drugs* 2001; **10**: 1409–1422.
93. Miller LA, Ratnam K, Payne DJ.  $\beta$ -Lactamase-inhibitor combinations in the 21st century: current agents and new developments. *Curr Opin Pharmacol* 2001; **1**: 451–458.
94. Koronakis V, Sharff A, Koronakis E, Luisi B, Hughes C. Crystal structure of the bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature* 2000; **405**: 914–919.
95. Murakami S, Nakashima R, Yamashita E, Yamaguchi A. Crystal structure of bacterial multidrug efflux transporter AcrB. *Nature* 2002; **419**: 587–593.
96. Avila-Sakar AJ, Misaghi S, Wilson-Kubalek EM *et al.* Lipid-layer crystallization and preliminary three-dimensional structural analysis of AcrA, the periplasmic component of a bacterial multidrug efflux pump. *J Struct Biol* 2001; **136**: 81–88.
97. Mao W, Warren MS, Black DS *et al.* On the mechanism of substrate specificity by resistance nodulation division (RND)-type multidrug resistance pumps: the large periplasmic loops of MexD from *Pseudomonas aeruginosa* are involved in substrate recognition. *Mol Microbiol* 2002; **46**: 889–901.
98. Tikhonova EB, Wang Q, Zgurskaya HI. Chimeric analysis of the multicomponent multidrug efflux trans-

- porters from Gram-negative bacteria. *J Bacteriol* 2002; **184**: 6499–6507.
99. Elkins CA, Nikaido H. Substrate specificity of the RND-type multidrug efflux pumps AcrB and AcrD of *Escherichia coli* is determined predominately by two large periplasmic loops. *J Bacteriol* 2002; **184**: 6490–6498.
100. Eda S, Maseda H, Nakae T. An elegant means of self-protection in Gram-negative bacteria by recognizing and extruding xenobiotics from the periplasmic space. *J Biol Chem* 2003; **278**: 2085–2088.
101. Yu EW, McDermott G, Zgurskaya HI, Nikaido H, Koshland DE. Structural basis of multiple drug-binding capacity of the AcrB multidrug efflux pump. *Science* 2003; **300**: 976–980.
102. Peng WT, Nester EW. Characterization of a putative RND-type efflux system in *Agrobacterium tumefaciens*. *Gene* 2001; **270**: 245–252.
103. Pumbwe L, Piddock LJ. Identification and molecular characterisation of CmeB, a *Campylobacter jejuni* multidrug efflux pump. *FEMS Microbiol Lett* 2002; **206**: 185–189.
104. Lin J, Michel LO, Zhang Q. CmeABC functions as a multidrug efflux system in *Campylobacter jejuni*. *Antimicrob Agents Chemother* 2002; **46**: 2124–2131.
105. Nagakubo S, Nishino K, Hirata T, Yamaguchi A. The putative response regulator BaeR stimulates multidrug resistance of *Escherichia coli* via a novel multidrug exporter system, MdtABC. *J Bacteriol* 2002; **184**: 4161–4167.
106. Baranova N, Nikaido H. The BaeSR two-component regulatory system activates transcription of the *yegM-NOB* (*mdtABCD*) transporter gene cluster in *Escherichia coli* and increases its resistance to novobiocin and deoxycholate. *J Bacteriol* 2002; **184**: 4168–4176.
107. Masuda N, Church GM. *Escherichia coli* gene expression responsive to levels of the response regulator EvgA. *J Bacteriol* 2002; **184**: 6225–6234.
108. Nishino K, Yamaguchi A. EvgA of the two-component signal transduction system modulates production of the YhiUV multidrug transporter in *Escherichia coli*. *J Bacteriol* 2002; **184**: 2319–2323.
109. Sulavik MC, Houseweart C, Cramer C *et al.* Antibiotic susceptibility profiles of *Escherichia coli* strains lacking multidrug efflux pump genes. *Antimicrob Agents Chemother* 2001; **45**: 1126–1136.
110. Ikeda T, Yoshimura F. A resistance-nodulation-cell division family xenobiotic efflux pump in an obligate anaerobe, *Porphyromonas gingivalis*. *Antimicrob Agents Chemother* 2002; **46**: 3257–3260.
111. Aendekerk S, Ghysels B, Cornelis P, Baysse C. Characterization of a new efflux pump, MexGHI–OpmD, from *Pseudomonas aeruginosa* that confers resistance to vanadium. *Microbiology* 2002; **148**: 2371–2381.
- 111a. Sekiya H, Mima T, Morita Y, Kuroda T, Mizushima T, Tsuchiya T. Functional cloning and characterization of a multidrug efflux pump, MexHI–OpmD, from a *Pseudomonas aeruginosa* mutant. *Antimicrob Agents Chemother* 2003; **47**: 2990–2992.
- 111b. Li Y, Mima T, Komori Y *et al.* A new member of the tripartite multidrug efflux pumps, MexVW–OprM, in *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 2003; **52**: 572–575.
112. Berlanga M, Vazquez JL, Hernandez-Borrell J, Montero MT, Vinas M. Evidence of an efflux pump in *Serratia marcescens*. *Microb Drug Resist* 2000; **6**: 111–117.
- 112a. Chen J, Kuroda T, Huda MN, Mizushima T, Tsuchiya T. An RND-type multidrug efflux pump SdeXY from *Serratia marcescens*. *J Antimicrob Chemother* 2003; **52**: 176–179.
113. Rahmati S, Yang S, Davidson AL, Zechiedrich EL. Control of the AcrAB multidrug efflux pump by quorum-sensing regulator SdiA. *Mol Microbiol* 2002; **43**: 677–685.
114. Wei Y, Lee JM, Smulski DR, LaRossa RA. Global impact of *sdiA* amplification revealed by comprehensive gene expression profiling of *Escherichia coli*. *J Bacteriol* 2001; **183**: 2265–2272.
115. Tsukagoshi N, Aono R. Entry into and release of solvents by *Escherichia coli* in an organic-aqueous two-liquid-phase system and substrate specificity of the AcrAB–TolC solvent-extruding pump. *J Bacteriol* 2000; **182**: 4803–4810.
116. Kopytek SJ, Dyer JC, Knapp GS, Hu JC. Resistance to methotrexate due to AcrAB-dependent export from *Escherichia coli*. *Antimicrob Agents Chemother* 2000; **44**: 3210–3212.
117. Jellen-Ritter AS, Kern WV. Enhanced expression of the multidrug efflux pumps AcrAB and AcrEF associated with insertion element transposition in *Escherichia coli* mutants selected with a fluoroquinolone. *Antimicrob Agents Chemother* 2001; **45**: 1467–1472.
- 117a. O'Neill AJ, Bostock JM, Morais MA, Chopra I. Antimicrobial activity and mechanisms of resistance to cephalosporin P1, an antibiotic related to fusidic acid. *J Antimicrob Chemother* 2002; **50**: 839–848.
118. Kobayashi K, Tsukagoshi N, Aono R. Suppression of hypersensitivity of *Escherichia coli* *acrB* mutant to organic solvents by integration activation of the *aceEF* operon with the IS 1 or IS2 element. *J Bacteriol* 2001; **183**: 2646–2453.
119. Veal WL, Shafer WM. Identification of a cell envelope protein (MtrF) involved in hydrophobic antimicrobial resistance in *Neisseria gonorrhoeae*. *J Antimicrob Chemother* 2003; **51**: 27–37.
120. Masuda N, Sakagawa E, Ohya S, Gotoh N, Tsujimoto H, Nishino T. Substrate specificities of MexAB–OprM, MexCD–OprJ, and MexXY–OprM efflux pumps in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2000; **44**: 3322–3327.
121. Okamoto K, Gotoh N, Nishino T. Alterations of susceptibility of *Pseudomonas aeruginosa* by overproduction of multidrug efflux systems, MexAB–OprM, MexCD–OprJ, and MexXY–OprM to carbapenems: substrate specificities of the efflux systems. *J Infect Chemother* 2002; **8**: 371–373.
122. Okamoto K, Gotoh N, Nishino T. Extrusion of penem antibiotics by multicomponent efflux systems MexAB–OprM, MexCD–OprJ, and MexXY–OprM of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2002; **46**: 2696–2699.
123. Koutsolioutsou A, Martins EA, White DG, Levy SB, Demple B. A *soxRS*-constitutive mutation contributing to antibiotic resistance in a clinical isolate of *Salmonella enterica* (serovar typhimurium). *Antimicrob Agents Chemother* 2001; **45**: 38–43.

124. Giraud E, Cloeckert A, Kerboeuf D, Chaslus-Dancla E. Evidence for active efflux as the primary mechanism of resistance to ciprofloxacin in *Salmonella enterica* serovar typhimurium. *Antimicrob Agents Chemother* 2000; **44**: 1223–1228.
125. Shyu JB, Lies DP, Newman DK. Protective role of *tolC* in efflux of the electron shuttle anthraquinone-2,6-disulfonate. *J Bacteriol* 2002; **184**: 1806–1810.
126. Sanchez P, Alonso A, Martinez JL. Cloning and characterization of *SmeT*, a repressor of the *Stenotrophomonas maltophilia* multidrug efflux pump *SmeDEF*. *Antimicrob Agents Chemother* 2002; **46**: 3386–3393.
127. Zhang L, Li X-Z, Poole K. The *SmeDEF* multidrug efflux pump contributes to intrinsic multidrug resistance in *Stenotrophomonas maltophilia*. *Antimicrob Agents Chemother* 2001; **45**: 3497–3503.
- 127a. Teran W, Felipe A, Segura A, Rojas A, Ramos JL, Gallegos M-T. Antibiotic-dependent induction of *Pseudomonas putida* DOT-T1E TtgABC efflux pump is mediated by the drug binding repressor TtgR. *Antimicrob Agents Chemother* 2003; **47**: 3067–3072.
- 127b. Rojas A, Segura A, Guazzaroni ME *et al.* In vivo and in vitro evidence that TtgV is the specific regulator of the TtgGHI multidrug and solvent efflux pump of *Pseudomonas putida*. *J Bacteriol* 2003; **185**: 4755–4763.
- 127c. Hirakawa H, Nishino K, Hirata T, Yamaguchi A. Comprehensive studies of drug resistance mediated by overexpression of response regulators of two-component signal transduction systems in *Escherichia coli*. *J Bacteriol* 2003; **185**: 1851–1856.
- 127d. Nishino K, Yamada J, Hirakawa H, Hirata T, Yamaguchi A. Roles of TolC-dependent multidrug transporters of *Escherichia coli* in resistance to  $\beta$ -lactams. *Antimicrob Agents Chemother* 2003; **47**: 3030–3033.
128. Miyamae S, Ueda O, Yoshimura F, Hwang J, Tanaka Y, Nikaido H. A MATE family multidrug efflux transporter pumps out fluoroquinolones in *Bacteroides thetaiotaomicron*. *Antimicrob Agents Chemother* 2001; **45**: 3341–3346.
129. Ninio S, Rotem D, Schuldiner S. Functional analysis of novel multidrug transporters from human pathogens. *J Biol Chem* 2001; **276**: 48250–48256.
130. Braibant M, Guilloteau L, Zygmunt MS. Functional characterization of *Brucella melitensis* NorMI, an efflux pump belonging to the multidrug and toxic compound extrusion family. *Antimicrob Agents Chemother* 2002; **46**: 3050–3053.
131. Wigfield SM, Rigg GP, Kavari M, Webb AK, Matthews RC, Burnie JP. Identification of an immunodominant drug efflux pump in *Burkholderia cepacia*. *J Antimicrob Chemother* 2002; **49**: 619–624.
132. Fehlner-Gardiner CC, Valvano MA. Cloning and characterization of the *Burkholderia vietnamiensis* *norM* gene encoding a multi-drug efflux protein. *FEMS Microbiol Lett* 2002; **215**: 279–283.
133. Rouquette-Loughlin C, Dunham SA, Kuhn M, Balthazar JT, Shafer WM. The NorM efflux pump of *Neisseria gonorrhoeae* and *Neisseria meningitidis* recognizes antimicrobial cationic compounds. *J Bacteriol* 2003; **185**: 1101–1106.
134. Li X-Z, Poole K, Nikaido H. Contributions of MexAB-OprM and an EmrE homologue to intrinsic resistance of *Pseudomonas aeruginosa* to aminoglycosides and dyes. *Antimicrob Agents Chemother* 2003; **47**: 27–33.
135. Chen J, Morita Y, Huda MN, Kuroda T, Mizushima T, Tsuchiya T. VmrA, a member of a novel class of Na (+)-coupled multidrug efflux pumps from *Vibrio parahaemolyticus*. *J Bacteriol* 2002; **184**: 572–576.